



Factsheet Heavy Metals



HEAVY METALS

Heavy metals are naturally occurring elements found throughout the earth's crust. However, human activities such as mining, industrial production and use, and domestic and agricultural use of metals and metal-containing compounds have caused heavy metals to distribute throughout the environment in the air, water, and land. Heavy metals are of public health significance as most are classified as human carcinogens and have the potential to cause multiple organ damage. The toxicity of heavy metals depends on several factors including the dosage, route of exposure, chemical form, and the age, gender, genetics, and nutritional status of the exposed individual. Exposure to heavy metals most often occurs through drinking contaminated water, eating contaminated food, using contaminated water in food preparation and irrigation of food crops, industrial processes, or smoking tobacco.

- Arsenic Arsenic is used as an alloying agent and in the processing of glass, pigments, textiles, paper, metal adhesives, wood preservatives and ammunition. Historically, and to a limited extent today, arsenic is used in pesticides. In certain geographical areas, there may be high levels of arsenic present in the soil where it was once used as a pesticide or in the groundwater because of leaching. Arsenic and its compounds are classified as Group 1 carcinogenic agents (carcinogenic to humans) by the International Agency for Research on Cancer.
- **Cadmium** Cadmium is used in the production of batteries, paint pigments, coatings and plating, stabilizers, and alloys. Cadmium sulphide is used as a pigment in plastics, ceramics, glasses, enamels, etc. Cadmium salts are used as stabilizers in PVC plastics and corrosion-resistant coatings for electronics, steel, and aluminium. Cadmium compounds are classified as Group 1 carcinogenic agents (carcinogenic to humans) by the International Agency for Research on Cancer.
- **Cobalt** Cobalt is used in the production of corrosion-resistant metal alloys, as a binding agent in cutting and drilling tools, and as a component in batteries, pigments, electronics, and medicine. Cobalt compounds are classified as Group 2B carcinogenic agents (possibly carcinogenic to humans) by the International Agency for Research on Cancer.
- **Copper** Copper is mainly used in construction and the manufacturing of electrical equipment and industrial machinery. Copper sulphate is a permitted food additive in wine as a fining agent. It can be added in such quantity that the content of copper in the finished product shall not exceed 1000 µg/L. Copper is also an essential element for biochemical and physiological functions in plants and animals. Copper is classified as a Group 3 carcinogenic agent (not classifiable as to its carcinogenicity to humans) by the International Agency for Research on Cancer.
- Lead Lead is mainly used in the production of lead-acid batteries for automobiles. Lead solder used in food cans has been virtually eliminated. Elemental lead is classified as a Group 2B carcinogenic agent (possibly carcinogenic to humans), inorganic lead compounds are classified as Group 2A carcinogenic agents (probably carcinogenic to humans), and organic lead compounds are classified as Group 3 carcinogenic agents (not classifiable as to its carcinogenicity to humans) by the International Agency for Research on Cancer.

HEAVY METAL LIMITS

Th	le fol	lowin	ng are	limits	establ	ished	by [.]	the	NSLC	for	the	heavy	metal	s outl	ined	abov	e:

Heavy Metal	Beer	Cider	Ready to Drink	Spirits	Wine	
Arsenic (µg/L)	Less than 100	Less than 100	Less than 100	Less than 100	Less than 100	
Cadmium (µg/L)	Less than 20	Less than 20	Less than 20	Less than 20	Less than 20	
Cobalt (µg/L)	Less than 20	Less than 20	Less than 20	Less than 20	Less than 20	
Copper (µg/L)	Not Tested	Not Tested	Not Tested	Not Tested	Less than 1000	
Lead (µg/L)	Less than 200	Less than 200	Less than 200	Less than 200	Less than 200	





SOURCE OF HEAVY METALS

Raw Materials

- Fruits, vegetables, grains, botanicals, etc. may take up heavy metals from soil and water through their roots. Heavy metals may also be deposited on raw materials from traffic fumes and other air pollution.
- Water used during the growing of crops may be contaminated with excess levels of heavy metals.
- Water used as an ingredient may be contaminated with excess levels of heavy metals.
- Although arsenic is generally no longer permitted in agricultural chemicals, it may still be present in the soil where it was previously used or have leached into groundwater increasing the content in the water supply.

During Processing

- Excess levels of copper sulphate used as a fining agent in wine. One study has shown that other fining agents, such as diatomaceous earth can contain elevated amounts of heavy metals.
- Water used during processing such as for washing bottles and equipment may be contaminated with excess levels of heavy metals.
- Different equipment and containers used during processing or storage may contain different heavy metals. For example, copper stills for whisky and other sources such as fermentation drums, pipes, barrels, and casks.

CONTROLLING HEAVY METAL CONTAMINATION

- Consider the raw material supplier. Ensure raw materials are purchased from approved, trusted suppliers. Consult with the supplier for further information on the presence of heavy metals in their raw materials, soil, and water supplies. Consider which pesticides are used and if the soil and water sources have been assessed for potential hazards. Visit Perennia's **website** for more information on commodity-specific production.
- Ensure all food additives, such as copper sulphate, are used in accordance with the maximum levels of use as outlined in <u>Health Canada's List of</u> <u>Permitted Additives</u>.
- Ensure water used as an ingredient and for processing and cleaning meets <u>Health Canada's Guidelines for</u> <u>Canadian Drinking Water Quality</u>. Heavy metals in water can be removed by treatment processes such as reverse osmosis. Regular testing of all water

sources used throughout production and processing is recommended. Maximum allowable concentrations and sources are as follows:

- » Arsenic 10µg/L or as low as reasonably achievable. Sources include naturally occurring (erosion and weathering of soils, minerals, ores), releases from mining, and industrial effluent.
- » Cadmium 7µg/L. Sources include leaching from galvanized pipes and solders, and industrial and municipal waste.
- » **Cobalt** Federal guidelines have not been established for cobalt in drinking water.
- » **Copper** 2000µg/L. Sources include naturally occurring and leaching from copper piping.
- » Lead 5µg/L or as low as reasonably achievable. Sources include leaching from plumbing (lead service lines, lead solder, and brass fittings).

REFERENCES

Agency for Toxic Substances and Disease Registry. (2011, October 24). ToxFAQsTM For Copper.

Carex Canada. (n.d.). Arsenic and Arsenic Compounds Profile.

Carex Canada. (n.d.). Cadmium Profile.

Carex Canada. (n.d.). Cobalt Profile.

Carex Canada. (n.d.). Lead Profile.

Donadini, G., Spalla, S., Beone, G. M. (2012). Arsenic, Cadmium and Lead in Beers from the Italian Market. Journal of the Institute of Brewing, 114(4): 283-288.

He, N. X., & Bayen, S. (2020). An overview of chemical contaminants and other undesirable chemicals in alcoholic beverages and strategies for analysis. Comprehensive Reviews in Food Science and Food Safety, 19(6): 3916-3950

Health Canada. (2021, October 20). Lead Information Package – Some Commonly Asked Questions About Lead and Human Health.

Ibanez, J. G., Carreon-Alvarez, A., Barcena-Soto, M., Casillas, N. (2008). Metals in alcoholic beverages: A review of sources, effects, concentrations, removal, speciation, and analysis. Journal of Food Composition and Analysis, 21(8): 672-683.

International Agency for Research on Cancer. (2010). IARC Monographs on the Evaluation of Carcinogenic Risks to Humans: Volume 96: Alcohol Consumption and Ethyl Carbamate. Lyon, France. International Agency for Research on Cancer.





Pang, X. N., Li, Z. J., Chen, J. Y., Gao, L. J., Han, B. Z. (2017). A Comprehensive Review of Spirit Drink Safety Standards and Regulations from an International Perspective. Journal of Food Protection, 80(3): 431-442.

Redan, B. W., Jablonski, J, E., Halverson, C., Jaganathan, J., Mabud, M. A., Jackson, L. S. (2019). Factors Affecting Transfer of the Heavy Metals Arsenic, Lead, and Cadmium from Diatomaceous-Earth Filter Aids to Acholic Beverages during Laboratory -Scale Filtration. Journal of Agricultural and Food Chemistry, 67(9): 2670-2678.

Royal Society of Chemistry. (n.d.) Copper.

Stockley, C., Paschke-Kratzin, A., Teissedre, P. L, Restani, P., Tejedor, N. G., Quini, C. (2021). Arsenic and Wine: A Review. International Organization of Vine and Wine Publications.

Tchounwou, P. B., Yedjou, C. G., Patlolla, A. K., Sutton, D. J. (2012). Heavy Metal Toxicity and the Environment. In Luch A. (eds) Molecular, Clinical and Environmental Toxicology. Experientia Supplementum, 101, 133–164. Springer, Basel.

United States Environmental Protection Agency. (2000, January). Cadmium Compounds.

United States Environmental Protection Agency. (2000, January). Cobalt Compounds.

United States Environmental Protection Agency (n.d.). Learn about Lead.

World Health Organization. (2018, February 15). Arsenic.

World Health Organization (2010). Exposure to Cadmium: A Major Public Health Concern.

FOR MORE INFORMATION

If you have questions about the information found in this factsheet, please contact one of Perennia's specialist at:

Quality and Food Safety

Phone: 902-896-0277 Email: foodsafety@perennia.ca

or

Food and Beverage Innovation Centre

Phone: 902-896-8782 Email: innovation@perennia.ca

If you have questions regarding the established limits or product testing, please contact the NSLC at **product.testing@mynslc.com**



